



## Rhythm Pharmaceuticals Announces Publication of Analysis of Quality of Life Improvements from Phase 3 Clinical Trial in Bardet-Biedl Syndrome

January 23, 2023

*- Patients reported clinically meaningful improvements across multiple health-related quality of life measures following 52 weeks of treatment with setmelanotide -*

*- Analysis published in Orphanet Journal of Rare Diseases -*

BOSTON, Jan. 23, 2023 (GLOBE NEWSWIRE) -- Rhythm Pharmaceuticals, Inc. (Nasdaq: RYTM), a commercial-stage biopharmaceutical company focused on transforming the lives of patients and their families living with hyperphagia and severe obesity caused by rare melanocortin-4 receptor (MC4R) pathway diseases, today announced that certain patients with Bardet-Biedl syndrome (BBS) and obesity who participated in the Company's global Phase 3 clinical trial of setmelanotide reported clinically meaningful improvements across multiple health-related quality of life (HRQOL) measures based on an analysis published in the *Orphanet Journal of Rare Diseases*.

"Living with hyperphagia and severe obesity associated with BBS presents a clinically significant burden on patients and families," said corresponding author Prof. Andrea M. Haqq, M.D., Department of Pediatrics, Faculty of Medicine & Dentistry, University of Alberta. "We are excited to share the results of this study that show adult, adolescent and pediatric patients with BBS achieved clinically meaningful improvements in HRQOL across physical function and psychosocial domains during this 52-week trial evaluating setmelanotide, further supporting setmelanotide as a targeted treatment for improving weight outcomes and quality of life for these patients."

In total, 20 out of 31 patients with BBS and obesity completed various self-reported HRQOL measures at baseline and after 52 weeks of treatment with setmelanotide, which is a precision medicine designed to re-establish MC4R pathway signaling that is approved by the U.S. Food & Drug Administration (FDA) for chronic weight management in adult and pediatric patients six years of age and older with monogenic or syndromic obesity due to BBS.

For the 52-week analysis, 9 patients younger than 18 years, or their caregiver, reported a clinically meaningful change in their quality of life with a mean change from baseline of +11.2 points on the Pediatric Quality of Life Inventory (PedsQL) measure. In addition, 11 adult patients who completed the Impact of Weight on Quality of Life Questionnaire-Lite (IWQOL-Lite) achieved a clinically meaningful improvement in their weight related quality of life with a mean change from baseline of +12.0 points. IWQOL-Lite score was significantly correlated with changes in percent body weight ( $P=0.0037$ ) and body mass index ( $P=0.0098$ ).

"The pathological, insatiable hunger and early-onset, severe obesity that many patients with BBS experience arises from impairment of the MC4R pathway and is particularly disruptive to quality of life for both patients and their families," said David Meeker, M.D., Chair, President and Chief Executive Officer of Rhythm. "We are focused on supporting patients with BBS and their families by delivering a precision medicine that can restore normal pathway signaling, and hope that by doing so we can help these patients experience better quality of life."

### **About Bardet-Biedl Syndrome**

Bardet-Biedl syndrome (BBS) is a heterogenous rare genetic syndromic disease that presents with a variety of symptoms that evolve over time including visual impairments, renal disease, polydactyly, genital abnormalities, cognitive impairment, and hyperphagia and early-onset, severe obesity arising from impairment of the hypothalamic MC4R pathway. In the United States, BBS affects approximately 2,500 individuals with similar prevalence in Europe.

### **About Rhythm Pharmaceuticals**

Rhythm is a commercial-stage biopharmaceutical company committed to transforming the lives of patients and their families living with hyperphagia and severe obesity caused by rare melanocortin-4 receptor (MC4R) pathway diseases. Rhythm's precision medicine, setmelanotide, is approved by the U.S. Food and Drug Administration (FDA) for chronic weight management in adult and pediatric patients 6 years of age and older with monogenic or syndromic obesity due to pro-opiomelanocortin (POMC), proprotein convertase subtilisin/kexin type 1 (PCSK1) or leptin receptor (LEPR) deficiency confirmed by genetic testing, or patients with a clinical diagnosis of Bardet-Biedl syndrome (BBS). The European Commission (EC) has authorized setmelanotide for the treatment of obesity and the control of hunger associated with genetically confirmed BBS or genetically confirmed loss-of-function biallelic POMC, including PCSK1, deficiency or biallelic LEPR deficiency in adults and children 6 years of age and above. The UK's Medicines & Healthcare Products Regulatory Agency (MHRA) authorized setmelanotide for the treatment of obesity and the control of hunger associated with genetically confirmed loss-of-function biallelic POMC, including PCSK1, deficiency or biallelic

LEPR deficiency in adults and children 6 years of age and above. Additionally, Rhythm is advancing a broad clinical development program for setmelanotide in other rare genetic diseases of obesity and is leveraging the Rhythm Engine and the largest known obesity DNA database -- now with approximately 45,000 sequencing samples -- to improve the understanding, diagnosis and care of people living with severe obesity due to certain genetic deficiencies. Rhythm's headquarters is in Boston, MA.

### **Setmelanotide Indication**

In the United States, setmelanotide is indicated for chronic weight management in adult and pediatric patients 6 years of age and older with monogenic or syndromic obesity due to POMC, PCSK1 or LEPR deficiency as determined by an FDA-approved test demonstrating variants in POMC, PCSK1 or LEPR genes that are interpreted as pathogenic, likely pathogenic, or of uncertain significance (VUS) or BBS.

In the European Union, setmelanotide is indicated for the treatment of obesity and the control of hunger associated with genetically confirmed Bardet-Biedl syndrome (BBS) or genetically confirmed loss-of-function biallelic pro-opiomelanocortin (POMC), including PCSK1, deficiency or biallelic leptin receptor (LEPR) deficiency in adults and children 6 years of age and above.

### **Limitations of Use**

In the United States and Europe, Setmelanotide should be prescribed and supervised by a physician with expertise in obesity with underlying genetic etiology.

Setmelanotide is not indicated for the treatment of patients with the following conditions as setmelanotide would not be expected to be effective:

- Obesity due to suspected POMC, PCSK1 or LEPR deficiency with *POMC*, *PCSK1* or *LEPR* variants classified as benign or likely benign
- Other types of obesity not related to POMC, PCSK1 or LEPR deficiency, or BBS, including obesity associated with other genetic syndromes and general (polygenic) obesity.

### **WARNINGS AND PRECAUTIONS**

**Skin Monitoring:** Setmelanotide may lead to generalized increased skin pigmentation and darkening of pre-existing naevi because of its pharmacologic effect. Full body skin examinations should be conducted annually to monitor pre-existing and new skin pigmentary lesions before and during treatment with setmelanotide.

**Heart rate and blood pressure monitoring:** Heart rate and blood pressure should be monitored as part of standard clinical practice at each medical visit (at least every 6 months) for patients treated with setmelanotide.

**Prolonged penile erection:** Spontaneous penile erections have been reported in clinical trials with setmelanotide. Patients who have a penile erection lasting longer than 4 hours should be instructed to seek emergency medical attention for potential treatment of priapism.

**Depression:** In clinical trials, depression has been reported in patients treated with setmelanotide. Patients with depression should be monitored at each medical visit during treatment with setmelanotide. Consideration should be given to discontinuing setmelanotide if patients experience suicidal thoughts or behaviors.

**Pediatric Population:** The prescribing physician should periodically assess response to setmelanotide therapy. In growing children, the impact of weight loss on growth and maturation should be evaluated. The prescribing physician should monitor growth (height and weight) using age- and sex-appropriate growth curves.

**Excipients:** This medicinal product contains 10 mg benzyl alcohol in each ml. Benzyl alcohol may cause allergic reactions. Patients who are pregnant or breastfeeding should be advised of the potential risk from the excipient benzyl alcohol, which might accumulate over time and cause metabolic acidosis. This medicinal product should be used with caution in patients with hepatic or renal impairment, because of the potential risk from the excipient benzyl alcohol which might accumulate over time and cause metabolic acidosis.

**Sodium:** This medicinal product contains less than 1 mmol sodium (23 mg) per dose, that is to say essentially "sodium-free."

### **ADVERSE REACTIONS**

The most frequent adverse reactions are hyperpigmentation (51%), injection site reaction (39%), nausea (33%), and headache (26%).

### **USE IN SPECIFIC POPULATIONS**

#### **Pregnancy**

There are no data from the use of setmelanotide in pregnant women. Animal studies do not indicate direct harmful effects with respect to reproductive toxicity. However, administration of setmelanotide to pregnant rabbits resulted in decreased maternal food consumption leading to embryo-foetal effects. As a precautionary measure, setmelanotide should not be started during pregnancy or while attempting to get pregnant as weight loss during pregnancy may result in fetal harm. If a patient who is taking setmelanotide has reached a stable weight and becomes pregnant, consideration should be given to maintaining setmelanotide

treatment as there was no proof of teratogenicity in the nonclinical data. If a patient who is taking setmelanotide and still losing weight gets pregnant, setmelanotide should either be discontinued, or the dose reduced while monitoring for the recommended weight gain during pregnancy. The treating physician should carefully monitor weight during pregnancy in a patient taking setmelanotide.

### **Breast-feeding**

It is unknown whether setmelanotide is excreted in human milk. A nonclinical study showed that setmelanotide is excreted in the milk of nursing rats. No quantifiable setmelanotide concentrations were detected in plasma from nursing pups. A risk to the newborn/infant cannot be excluded. A decision must be made whether to discontinue breastfeeding or to discontinue/abstain from setmelanotide therapy taking into account the benefit of breastfeeding for the child and the benefit of therapy for the mother.

### **Fertility**

No human data on the effect of setmelanotide on fertility are available. Animal studies did not indicate harmful effects with respect to fertility.

To report SUSPECTED ADVERSE REACTIONS, contact Rhythm Pharmaceuticals at +1 (833) 789-6337. See [Summary of Product Characteristics' APPENDIX V](#) for a list of European national reporting systems to communicate adverse reactions.

**Please see the full Prescribing Information for additional Important Safety Information.**

### **Forward-Looking Statements**

This press release contains forward-looking statements within the meaning of the U.S. Private Securities Litigation Reform Act of 1995. All statements contained in this press release that do not relate to matters of historical fact should be considered forward-looking statements, including without limitation statements regarding the potential, safety, efficacy, and regulatory and clinical progress of setmelanotide, the potential benefits of setmelanotide for patients, including those with Bardet-Biedl Syndrome, and our business strategy and plans, including regarding commercialization of setmelanotide. Statements using word such as “expect”, “anticipate”, “believe”, “may”, “will” and similar terms are also forward-looking statements. Such statements are subject to numerous risks and uncertainties, including, but not limited to, our ability to enroll patients in clinical trials, the design and outcome of clinical trials, the impact of competition, the ability to achieve or obtain necessary regulatory approvals, risks associated with data analysis and reporting, our liquidity and expenses, the impact of the COVID-19 pandemic on our business and operations, including our preclinical studies, clinical trials and commercialization prospects, and general economic conditions, and the other important factors discussed under the caption “Risk Factors” in our Quarterly Report on Form 10-Q for the quarter ended September 30, 2022 and our other filings with the U.S. Securities and Exchange Commission. Except as required by law, we undertake no obligations to make any revisions to the forward-looking statements contained in this release or to update them to reflect events or circumstances occurring after the date of this release, whether as a result of new information, future developments or otherwise.

### **Corporate Contact:**

David Connolly  
Head of Investor Relations and Corporate Communications  
Rhythm Pharmaceuticals, Inc.  
857-264-4280  
[dconnolly@rhythmtx.com](mailto:dconnolly@rhythmtx.com)

### **Investor Contact:**

Hannah Deresiewicz  
Stern Investor Relations, Inc.  
212-362-1200  
[hannah.deresiewicz@sternir.com](mailto:hannah.deresiewicz@sternir.com)

### **Media Contact:**

Adam Daley  
Berry & Company Public Relations  
212-253-8881  
[adaley@berrypr.com](mailto:adaley@berrypr.com)



Source: Rhythm Pharmaceuticals, Inc.